JCI The Journal of Clinical Investigation

RANKL inhibition improves muscle strength and insulin sensitivity and restores bone mass

Nicolas Bonnet, ..., Eleni Douni, Serge Ferrari

J Clin Invest. 2023;133(4):e169317. https://doi.org/10.1172/JCI169317.

Corrigendum

Original citation: J Clin Invest. 2019;129(8):3214–3223. https://doi.org/10.1172/JCI125915 Citation for this corrigendum: J Clin Invest. 2023;133(4):169317. https://doi.org/10.1172/JCI169317 The authors recently became aware that an incorrect image was shown in Figure 6F for the Pparb–/– Opg-Fc sample. The correct figure is shown below. The authors regret the error.

Find the latest version:



Corrigendum

RANKL inhibition improves muscle strength and insulin sensitivity and restores bone mass

Nicolas Bonnet, Lucie Bourgoin, Emmanuel Biver, Eleni Douni, and Serge Ferrari

Original citation: *J Clin Invest*. 2019;129(8):3214–3223. https://doi.org/10.1172/JCI125915.

Citation for this corrigendum: J Clin Invest. 2023;133(4):169317. https://doi.org/10.1172/JCI169317.

The authors recently became aware that an incorrect image was shown in Figure 6F for the *Pparb*-/- Opg-Fc sample. The correct figure is shown below.

The authors regret the error.



Figure 6. Bone, muscle, and glucose phenotype of *Pparb*^{-/-} mice treated by OPG-Fc. (A and B) Skeletal muscle volume of the limb and fat infiltration in muscle evaluated by in vivo microCT. (C) Maximal speed evaluated on treadmill normalized by gastrocnemius mass. (D) Limb force evaluated by handgrip normalized by gastrocnemius mass (n = 8 per group). (E) Body temperature evaluated by infrared camera (n = 8 per group). (F) Muscle fiber type, number, and area. Note the type I fibers in blue dark and type II fibers in light blue. Original magnification is ×10. (G) ITT AUC (n = 8 per group). (H) GTT. (I) Relative protein expression in the gastrocnemius. Hatch marks correspond to mice that have received an acute injection of insulin. (J) Relative mRNA expression of insulin signaling in soleus. (K) Relative mRNA expression of Nfkb signaling in soleus (n = 6 per group). Statistical differences were assessed by 1-way ANOVA.